



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2017

Echocardiography accurately predicts pulmonary hypertension in patients with advanced lung disease

Cottini, Silvia ; Benden, Christian ; Huber, Lars C ; Arrigo, Mattia

DOI: <https://doi.org/10.1186/s13054-017-1697-y>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-143960>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Cottini, Silvia; Benden, Christian; Huber, Lars C; Arrigo, Mattia (2017). Echocardiography accurately predicts pulmonary hypertension in patients with advanced lung disease. *Critical Care*, 21(1):115.

DOI: <https://doi.org/10.1186/s13054-017-1697-y>

LETTER

Open Access



Echocardiography accurately predicts pulmonary hypertension in patients with advanced lung disease

Silvia Cottini¹, Christian Benden², Lars C. Huber^{2,3} and Mattia Arrigo^{4*} 

Keywords: Pulmonary hypertension, Echocardiography, Advanced lung disease, Transplantation, Right heart catheterization

Pulmonary hypertension (PH) may be observed in many different conditions, including advanced heart or lung disease [1]. PH is defined as elevation of the mean pulmonary arterial pressure (mPAP) ≥ 25 mmHg, measured by right heart catheterization (RHC). Since procedure-related risks and costs are not negligible, RHC is not routinely performed in patients with advanced lung disease during initial assessment and follow-up. Echocardiography is commonly used as screening tool for the presence of PH, but might be of limited diagnostic value, in particular in patients with lung disease [2]. Empirical formulas were proposed to calculate the mPAP starting from systolic pulmonary arterial pressure (sPAP) as assessed by echocardiography [3, 4]. It has been suggested that both the Chemla formula ($\text{mPAP} = \text{sPAP} \times 0.61 + 2$ mmHg) and the Syed formula ($\text{mPAP} = \text{sPAP} \times 0.65 + 0.55$ mmHg) might accurately estimate the mPAP [5]. The aim of our study was to assess the accuracy of these echocardiography-based formulas to estimate the invasively measured mPAP in a cohort of patients with advanced lung disease. Estimation of sPAP by echocardiography was performed according to current recommendation by adding to the right ventricular/right atrial pressure gradient (based on Doppler-measured tricuspid regurgitation velocity) the estimated right atrial pressure (based on inferior vena cava diameter and collapsibility). A total

of 96 consecutive patients undergoing lung transplant evaluation between 03/2000 and 10/2015 were included. Data from RHC and echocardiography performed the same day were available for the whole cohort.

Spearman's rank-order correlation was run to assess the relationship between RHC-measured mPAP and calculated mPAP. There was a strong positive correlation between both RHC-mPAP and Chemla-mPAP ($r_s = 0.909$, $p < 0.001$; Fig. 1a) and between RHC-mPAP and Syed-mPAP ($r_s = 0.904$, $p < 0.001$; Fig. 1b).

Inspection of both diagrams suggests that the correlation between echocardiographic data and invasive pulmonary hemodynamics is of particular strength in patients with moderate elevation of pulmonary pressure. Whether echocardiography-based formulas are of similar accuracy to estimate pulmonary pressure of PH in patients with severe PH remains unclear. However, this might be of minor relevance since, as shown in our cohort, only a minority of patients with advanced lung disease present with a mPAP > 50 mmHg. In these patients, further invasive assessment of pulmonary hemodynamics may be indicated. For all other patients our data emphasize that echocardiography is an accurate tool to estimate pulmonary pressure in patients with advanced lung disease.

* Correspondence: mattia.arrigo@usz.ch

⁴Department of Cardiology, University Hospital Zurich, Raemistrasse 100, 8091 Zurich, Switzerland

Full list of author information is available at the end of the article



© The Author(s). 2017 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.




Fig. 1 Correlation between RHC-measured mPAP and echocardiographic-estimated mPAP using the Chemla formula (a) and the Syed formula (b). Values expressed as mmHg

Abbreviations

mPAP: Mean pulmonary arterial pressure; PH: Pulmonary hypertension;
RHC: Right heart catheterization; sPAP: Systolic pulmonary arterial pressure

Acknowledgements

Not applicable.

Funding

Departmental funds.

Availability of supporting data

All supporting data are available upon request.

Authors' contributions

SC participated in the design of the study, collected the data, and drafted the manuscript. CB participated in the design of the study and critically revised the manuscript. LCH participated in the design of the study and drafted and revised the manuscript. MA performed statistical analyses and drafted and revised the manuscript. All authors gave final approval of the manuscript. All authors participated in data analysis and manuscript draft and revision. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Publication was authorized by the local Ethical Committee (BASEC 2016-01642).

Ethics approval and consent to participate

The study was performed according to the principles of the Declaration of Helsinki and was authorized by the local Ethical Committee (Kantonale Ethikkommission Zürich, Switzerland: BASEC 2016-01642).

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Intensive Care Unit, University Hospital Zurich, Zurich, Switzerland. ²Division of Pulmonology, University Hospital Zurich, Zurich, Switzerland. ³Clinic for Internal Medicine, City Hospital Triemli, Zurich, Switzerland. ⁴Department of Cardiology, University Hospital Zurich, Raemistrasse 100, 8091 Zurich, Switzerland.

Published online: 25 May 2017

References

1. Arrigo M, Huber LC. Passive pulmonary hypertension: more than hydrostatics. *Chest*. 2014;145:413.
2. Janda S, Shahidi N, Gin K, Swiston J. Diagnostic accuracy of echocardiography for pulmonary hypertension: a systematic review and meta-analysis. *Heart*. 2011;97:612–22. <http://heart.bmj.com/cgi/doi/10.1136/hrt.2010.212084>.
3. Chemla D, Castelain V, Provencher S, Humbert M, Simonneau G, Hervé P. Evaluation of various empirical formulas for estimating mean pulmonary artery pressure by using systolic pulmonary artery pressure in adults. *Chest*. 2009;135:760–8.
4. Syed R, Reeves JT, Welsh D, Raeside D, Johnson MK, Peacock AJ. The relationship between the components of pulmonary artery pressure remains constant under all conditions in both health and disease. *Chest*. 2008;133:633–9.
5. Aduen JF, Castello R, Lozano MM, Hepler GN, Keller CA, Alvarez F, et al. An alternative echocardiographic method to estimate mean pulmonary artery pressure: diagnostic and clinical implications. *J Am Soc Echocardiogr*. 2009; 22:814–9.